

## CHECK-LIST OF GNATHOSTOMULIDS FROM THE CANARY ARCHIPELAGO (NE ATLANTIC OCEAN)

**Rodrigo Riera\***

Centro de Investigaciones Medioambientales del Atlántico  
Arzobispo Elías Yanes 44, 38206 La Laguna, Tenerife, Canary Islands, Spain  
\*email: rodrigo@cimacanarias.com

### RESUMEN

Un total de cinco especies de gnatostomúlidos han sido citadas para el archipiélago canario. Las especies son: *Haplognathia rosea* (Sterrer, 1969), *H. ruberrima* (Sterrer, 1966), *Paragnathiella trifoliceps* Sterrer, 1997, *Labidognathia longicollis* Riedl, 1970 y *Austrognathia clavigera* Sterrer, 1997. Dos especies son endémicas (*Paragnathiella trifoliceps* y *Austrognathia clavigera*) y únicamente han sido recolectadas en su localidad tipo. El resto de especies se caracterizaron por presentar una distribución anfiatlántica (*L. longicollis*) y cosmopolita (*H. rosea* y *H. ruberrima*).

**Palabras clave:** Gnathostomulida, intersticial, sedimento, submareal, islas Canarias, Océano Atlántico

### ABSTRACT

Five species of gnathostomulids have been recorded from the Canary archipelago. These species are *Haplognathia rosea* (Sterrer, 1969), *H. ruberrima* (Sterrer, 1966), *Paragnathiella trifoliceps* Sterrer, 1997, *Labidognathia longicollis* Riedl, 1970 and *Austrognathia clavigera* Sterrer, 1997. Two species are endemisms, only recorded in type locality (*P. trifoliceps* and *A. clavigera*). The remaining taxa are amphiatlantic (*L. longicollis*) and cosmopolitan (*H. rosea* and *H. ruberrima*).

**Key words:** Gnathostomulida, interstitial, sediment, subtidal, Canary Islands, Atlantic Ocean.

### INTRODUCTION

Gnathostomulids are interstitial organisms, with an average length of 1.5 mm and a diameter ranging from 0.045 to 0.065 mm. Most of gnathostomulids live in marine anoxic sandy sediments, being common inhabitants of the RPD layer (REISE [4]). Up to now, almost 100 species of this taxonomic group has been described worldwide (STERRER [12]).

In Canary Islands, STERRER [11] found several gnathostomulid species in Las Canteras beach (Gran Canaria) in February 1996, even he discovered two new species (*Paragnathiella trifoliceps* and *Austrognathia clavigera*), one of them belonging to a new genus (*Paragnathiella*). Posteriorly, during a pilot study prior of one-year study basis conducted in Los Cristianos beach, several specimens belonging to two gnathostomulid species were recorded. A taxonomic detailed study revealed that they correspond to previously collected species by STERRER [11] in Gran Canaria: *Haplognathia rosea* and *H. ruberrima*.

## MATERIAL AND METHODS

Sediment samples were collected by hand and snorkelling, and extracted following methodology explained in STERRER [9, 10]. Samples from shallow subtidal seabeds of Los Cristianos beach were taken following methods described in detail by RIERA [6].

## SYSTEMATICS

**PHYLUM GNATHOSTOMULIDA** Ax, 1956

**Order FILOSPERMOIDEA** Sterrer, 1972

**Family Haplognathiidae** Sterrer, 1972

Genus *Haplognathia* Sterrer, 1970

*Haplognathia rosea* (Sterrer, 1969)

*Pterognathia rosea* STERRER [8]

*Haplognathia rosea* STERRER [11]: 186, Figs 1A, 2A.

**Studied material.**- Los Cristianos bay (Tenerife), coordinates (28°02'67"N, 16°42'64"W), 4 m depth, fine sands with very sparse *Cymodocea nodosa* patches, April 2000, 2 ind.

**Distribution.**- Cosmopolitan, collected in North Sea (STERRER [8]), Atlantic Ocean (STERRER [11]) and Pacific Ocean (STERRER [10]). This species was recorded in two sites in Las Canteras beach in shallow seabeds (2-3 m depth) with sparse *C. nodosa* patches and. rhizomes (STERRER [11]).

*Haplognathia ruberrima* (Sterrer, 1966)

*Pterognathia ruberrima* STERRER [7]

*Pterognathia grandis* KIRSTEUER [3]

*Haplognathia ruberrima* STERRER [11]: 186, Figs: 1B-G, 2B-C; Table 2.

**Studied material.**- Los Cristianos bay (Tenerife), coordinates (28°02'67"N, 16°42'64"W), 4 m depth, fine sands with very sparse *Cymodocea nodosa* patches, April 2000, 2 ind.

**Distribution.**- Cosmopolitan. North Sea (STERRER [8]), Atlantic Ocean (STERRER [11]) and Pacific Ocean (STERRER [10]). This species was recorded in Las Canteras beach, at shallow seabed (2-3 m depth) fine sands with sparse *C. nodosa* (STERRER [11]).

**Order BURSOVAGINOIDEA** Sterrer, 1972  
**Suborder Scleroperalia** Sterrer, 1972  
**Family Agnathiellidae** Sterrer, 1972  
Genus *Paragnathiella* Sterrer, 1997  
*Paragnathiella trifoliceps* Sterrer, 1997

*Paragnathiella trifoliceps* STERRER [11]: 189, Figs. 3A-L, 4A-E; Table 3.

**Distribution.-** Canary Islands. The type locality is Las Canteras beach, in fine sands with sparse *Cymodocea nodosa* at 2-3 m depth (STERRER [11]).

**Family Mesognathariidae** Sterrer, 1972  
Genus *Labidognathia* Riedl, 1970  
*Labidognathia longicollis* Riedl, 1970

*Labidognathia longicollis* RIEDL [5]: 229, fig. 1; STERRER [11]: 193, Figs. 5A-E, 6A-B; Table 4.

**Distribution.-** Amphiatlantic (RIEDL [5], STERRER [11]). This species has been recorded at Las Canteras beach, in fine sands with sparse *Cymodocea nodosa* at 2-3 m depth (STERRER [11]).

**Suborder CONOPHORALIA** Sterrer, 1972  
**Family Austrognathiidae** Sterrer, 1971  
Genus *Austrognathia* Sterrer, 1965  
*Austrognathia clavigera* Sterrer, 1997

*Austrognathia clavigera* STERRER [11]: 194, Figs. 7A-O, 8A-C, Table 5.

**Distribution.-** Canary Islands. The type locality is Las Canteras beach, in fine sands with sparse *Cymodocea nodosa* at 2-3 m depth (STERRER [11]).

## DISCUSSION

Most of taxonomic groups from interstitial fauna, commonly known as meiofauna, remains scarcely known in the Canary archipelago and have not been extensively studied, such as, gastrotrichs, turbellarians, harpacticoid copepods, acari, oligochaetes, priapulids, kinorhynchs, among others. Though several taxonomic works on free-living marine nematodes and polychaetes has been published from two sampling locations on the south coast of Tenerife (Los Abrigos del Porís and Los Cristianos) in the last decade (RIERA [6]) and several sea-grass meadows throughout the Canary archipelago (BRITO-CASTRO [2]), as well as, taxonomic studies carried out in the benthos laboratory of the University of La Laguna (Department of Animal Biology) coordinated by Dr. Jorge Núñez, most of the meiofaunal groups have only recorded in a limited number of sampling stations conducted by taxonomists during non-intensive field surveys. Hence, taxonomic extensive studies are necessary to carry out in order to increase the knowledge of canarian marine biodiversity.

Gnathostomulids are one of the interstitial taxonomic groups that still remain overlooked in the Canary archipelago. An increase number of gnathostomulids records is expected

in the next years, though specific sampling campaigns are necessary to be conducted because of ecologic requirements of this group (e.g. hypoxic or anoxic sandy sediments) (REISE [4]).

Current assemblages of gnathostomulid endemisms (*Paragnathiella trifoliceps* and *Austrognathia clavigera*) could be considered endangered, since they were recorded solely in the sparse *Cymodocea nodosa* seagrass meadow at Las Canteras beach (Gran Canaria). In the last years, a dramatic reduction of *C. nodosa* meadow has been observed; even demonstrated by genetic analyses that confirmed higher fragmentation and smaller meadow size in this location (ALBERTO *et al.* [1]).

## ACKNOWLEDGEMENTS

To Viceconsejería de Medio Ambiente del Gobierno de Canarias for coordinating BIOTA MARINO programme. To Leopoldo Moro and Rogelio Herrera (Canarian Government) for their continuous encouragement in the study of canarian marine biodiversity.

## REFERENCES

- [1] ALBERTO, F., S. ARNAUD-HAOND, C.M. DUARTE & E.A. SERRÃO. 2006. Genetic diversity of a clonal angiosperm near its range limit: the case of *Cymodocea nodosa* at the Canary Islands. *Marine Ecology Progress Series*, 309: 117-129.
- [2] BRITO-CASTRO, M.C. 2002. *Estudio de las comunidades intersticiales del sebadal (Cymodocea nodosa) en Canarias, con especial referencia a anélidos poliquetos*. Tesis Doctoral, curso 1998-1999. Servicio de Publicaciones de la Universidad de La Laguna.
- [3] KIRSTEUEER, E. 1969. On some species of Gnathostomulida from Bimini, Bahamas. *Amer. Mus. Novitates*, 2356: 1-21.
- [4] REISE, K. 1981. Gnathostomulida abundant alongside polychaete burrows. *Marine Ecology Progress Series* 6: 329-333.
- [5] RIELD, R. 1970. On *Labidognathia longicollis* nov. gen. spec. from the West Atlantic Coast (Gnathostomulida). *Int. Revue ges. Hydrobiol.* 55: 359-376.
- [6] RIERA, R. 2004. *Biodiversidad meiofaunal de las playas de Los Abrigos del Porís y de Los Cristianos en la isla de Tenerife. Estructura y dinámica de sus comunidades*. University of La Laguna, Ph-D Thesis, 486 pp.
- [7] STERRER, W. 1966. *Gnathostoula paradoxa* und vertreter von *Pterognathia* (ein neues Gnathostomuliden-Genus) von der schwedischen Westküste. *Arkiv för Zoologi*, Serie 2, 18(16): 405-413.
- [8] STERRER, W. 1969. Beiträge zur Kenntnis der Gnathostomulida I. Anatomie und Morphologie des Genus *Pterognathia* Sterrer. *Arkiv för Zoologi*, Serie 2, 22(1): 1-125.
- [9] STERRER, W. 1971. On the biology of Gnathostomulida. *Vie et Milieu*, suppl 22: 493-508.
- [10] STERRER, W. 1991. Gnathostomulida from Fiji, Tonga and New Zealand. *Zoologica Scripta*, 20: 107-128.
- [11] STERRER, W. 1997. Gnathostomulida from the Canary Islands. *Proceedings of the Biological Society of Washington*, 110(2): 186-197.
- [12] STERRER, W. 2006. Gnathostomulida (unsegmented marine worms). In eLS. John Wiley & Sons Ltd, Chichester.